

# ASSESSING THE MATERNAL AND FETAL OUTCOMES AMONG THE ECLAMPSIA PATIENTS ATTENDING AT DHAKA MEDICAL COLLEGE HOSPITAL OF BANGLADESH: A CROSS-SECTIONAL STUDY

Farida Iasmin<sup>1,3</sup>, Ayesha Akter<sup>1,2</sup>, Tasnim Akter<sup>1,4</sup>, ARM Myeenuddin Chowdhury<sup>5</sup>

<sup>1</sup>Department of Public Health, Daffodil International University, Dhaka, Bangladesh.

<sup>2</sup>HBH International Nursing College, Sector #10, Uttara, Dhaka 1230, Bangladesh.

<sup>3</sup>National Institute of Diseases of the Chest and Hospital, Mohakhali, Dhaka-1212, Bangladesh.

<sup>4</sup>East West Nursing College, Turag, Dhaka 1711, Bangladesh.

<sup>5</sup>Faculty of Environmental Earth Science, Hokkaido University, Hokkaido, Japan

## Abstract

*Eclampsia is a life-threatening emergency that contributes to major causes of serious maternal morbidity and mortality worldwide. This descriptive type of cross-sectional study was conducted among 125 eclampsia patients at Eclampsia ward in Dhaka Medical College and Hospital from January to August 2015. The study was aimed to assess the maternal and fetal outcomes of eclampsia patients in Bangladesh. Data were collected through face to face interview by using a pretested semi-structure questionnaire. Study results showed that majority of the respondents' age was between 20-25 years and a number of respondents (66%) had caesarean delivery. Pulmonary edema, HELLP syndrome, obstetric shock was found as maternal complications due to eclampsia and maternal death rate was 0.8%. This study also found that perinatal death due to eclampsia was 8.0% and 6.4% were stillborn and 12.8% were IUD. Birth asphyxia, low birth weight, IUGR, prematurity was evident as important causes of neonatal death among eclamptic mothers.*

**Keywords:** Eclampsia, Maternal outcome, Foetal outcome, Public health, Bangladesh.

## Introduction

Eclampsia is a life-threatening obstetric complication characterized by hypertension, proteinuria and convulsion with unknown etiology<sup>1</sup>. It is the occurrence of convulsion in association with the feature of pre-eclampsia which requires immediate diagnosis and management<sup>2</sup>. Eclampsia is a multisystem disorder that involves failure of vital organs which may leads to deterioration of maternal and foetal condition resulting in high maternal and perinatal mortality and morbidity<sup>3</sup>.

---

\*Corresponding Author: Tasnim Akter, Department of Public Health, Daffodil International University (DIU), Bangladesh. Email: [tasnimmunni23@gmail.com](mailto:tasnimmunni23@gmail.com)

Preeclampsia and eclampsia are major causes of maternal and perinatal morbidity and mortality. Transient neurological deficit is common but persistent deficits are rare<sup>4</sup>. Perinatal mortality rates of up to 19.1% are reported from centers in Nigeria in association with eclampsia<sup>5</sup>. The incidence varies widely in different countries and even in the different zones of the same country. The incidence of eclampsia in developed countries approximately 1 in 2000 deliveries<sup>6</sup>. In developing countries, it is from 1 in 100 to 1 in 1700<sup>7, 8</sup>. More than 5 lac women die of pregnancy related causes in each year and 99% deaths occur in the developing countries<sup>9, 10</sup>. Among them 50,000 maternal deaths occur due to eclampsia in every year<sup>11</sup>.

In Bangladesh, about 4500 women die due to eclampsia which contributes 16% of the maternal mortality in one year<sup>12</sup>. At present, the perinatal mortality rate in Bangladesh is 65 per 1,000 live births<sup>13</sup>. Eclampsia is associated with increased rates of perinatal mortality and morbidity. Two different studies in Bangladesh showed perinatal mortality rate due to Eclampsia were 28% and 32%<sup>14, 15</sup>. The cause behind the high incidence of maternal and perinatal mortality and morbidity are lack of antenatal care, low socio-economic status, illiteracy, ignorant of disease, substandard care, shyness and religious belief etc. Besides poor communication facilities and social superstition are also important factors<sup>16</sup>.

There are approximately 3.6 million births per year in Bangladesh and over 10,000 women develop eclampsia each year<sup>17</sup>. It is one of the common causes of maternal mortality and responsible for 20 % of maternal death<sup>18</sup>. Low educational status and low socioeconomic condition among the women of Bangladesh is the main cause of absence of antenatal care and development of eclampsia. Eclampsia is a major cause of stillbirth and neonatal death, intra uterine growth retardation (IUGR), low birth weight (LBW), prematurity and neonatal asphyxia are another consequence of eclampsia. Though a number of patients delivered a healthy baby with good outcome, perinatal mortality is high to the extent of 30-50 percent in eclampsia<sup>19</sup>.

A study on factors associated with severe preeclampsia and eclampsia in Jahun, Nigeria was found that 16% women had eclampsia who had history of previous preeclampsia and hypertension<sup>20</sup>. Maternal and perinatal case fatality rates were 7.89% and 20.73% respectively among eclamptic patients in Tanzania<sup>21</sup>. Another study conducted in Nepal was also found that eclampsia is associated with poor fetomaternal outcome<sup>22</sup>. Maternal complications like pulmonary edema, intra-cerebral hemorrhage, disseminated intravascular coagulopathy (DIC), acute renal failures etc. are serious reasons causing maternal mortality and morbidity<sup>25, 26, 27</sup>. The most common cause of foetal deaths are pre-maturity and birth asphyxia <sup>[28]</sup>. A previous study was assessed knowledge of pre-eclampsia and eclampsia among women in Bangladesh<sup>23</sup>. Another prospective study was done in Chuadanga Sadar Hospital, to measure the outcome of eclampsia patients in district hospital in Bangladesh<sup>24</sup>. Research conducted in Sylhet region of Bangladesh was identified the prevalence of preeclampsia and the associated risk factors among the

pregnant women<sup>36</sup>. However, there is limited of data regarding maternal and fetal outcomes among the Eclampsia patients attending at Dhaka Medical College Hospital of Bangladesh. As a consequence, this study was designed to assess the maternal and fetal outcomes among the Eclampsia patients attending at Dhaka Medical College Hospital of Bangladesh. This study may provide some guidelines for establishing plans to improve maternal and perinatal outcomes and could generate interest in further study.

## **Materials & methods**

### **Study design & settings**

This descriptive type of cross-sectional study was conducted in the department of Dhaka Medical College Hospital (DMCH) located in the Dhaka city of Bangladesh. The study was conducted during the time period of January to August 2021.

### **Study population, sample size and sampling strategy**

Study population was the eclampsia patients admitted at Dhaka Medical College Hospital (DMCH) during data collection period. A total of 125 samples were taken for data collection. Purposive non-probability sampling technique was used to select the target sample size. Eclampsia patients who were severely ill, non-cooperative and unwilling to participate were excluded from the study.

### **Data collection procedure and quality control**

Data was gathered by using a semi-structured interview questionnaire. The questionnaire's suitability, appropriateness, acceptability, and sequences of the questions were pretested among 15 members of the eclamptic patients who were not sampled. Face-to-face interviews were done with respondents for data collection. Before starting the interview a written informed consent was taken from the respondents and ensured they understood the study's objectives.

The fact that the respondents might choose whether or not to respond to any question was made clear to them. Regarding some ethical issues, they were given complete assurance that the study's findings would never be disclosed to anyone who wasn't participating in it or used for the purpose for which it was intended.

### **Data analysis**

After interviews, each questionnaire was carefully reviewed for accuracy, completeness, and internal consistency to make sure no information was left missing or inconsistent. Prior to coding and analysis, the researchers themselves evaluated and edited all of the data using "Statistical Package for Social Science (SPSS) version 16.0." The analytical strategy was developed while keeping in mind the study's objectives. The data was presented by using tables, graphs, and charts.

### **Ethical Issue**

The study proposal was submitted to 'FAHS Research Ethics Committee, Daffodil International University or approval. Before collecting data, the study volunteers were taken written informed consent from the participants. The participants identities were protected, and they were made aware that they could withdraw at any time during the data collection process. Anonymity and confidentiality was strictly maintained. Administrative approval was also taken from the Dhaka Medical College Hospital authority.

### **Results**


#### **Socio-demographic information of the participants**

**Table-1** represents the socio-demographic characteristics of the respondent. A total of 125 Eclampsia patients were interviewed. Out of 125 respondents 21(16.8%) of the respondent are 15-19 years of age, 56(44.8%) respondent are 20-25 years, 33(26.4%) respondent are 26-30 years, 15(12.0%) respondent are >30 years. The monthly family income of the majority of respondents 54(43.2%) were between 10,001-15,000 TK, then 27(21.6%) earned <10,000 Tk, and similar 22(17.6%) earned 15,001-20,000 Tk, 22(17.6%) earned >20,000 Tk. Maximum respondents 68(54.4%) were lived in rural areas whereas remaining 57(45.6%) were lived in urban areas. Among 125 respondents 59(47.2%) had completed primary education, 37(29.6%) had secondary school, only 16(12.8%) had completed above secondary school, and 13(10.4%) respondents were informal education. Among this study group majority of respondents 108(86.4%) were housewife, 6(4.8%) were service holder, 1(0.8%) had business and 10(8.0%) were others service. 56(44.8%) respondents husband completed primary education, 44(35.2%) husband completed secondary school, 24(19.2%) husband had above secondary school, only 1(0.8%) husband had informal education. Among this study group majority of respondents husband 53(42.4%) were others service, 34(27.2%) were business, 24(19.2%) were service holder and 14(11.2%) were farmer (**Table1**).

**Table 1 Socio-demographic characteristics of the participants**


<b>Participants' characteristics</b>		<b>Frequency</b>	<b>%</b>
<b>Age</b>	15-19 years	21	16.8
	20-25 years	56	44.8
	26-30 years	33	26.4
	>30 years	15	12.0
	<b>Mean ± SD</b>	<b>24.45 ± 5.2</b>	
<b>Monthly family income</b>	< 10,000 Tk	27	21.6
	10,001-15,000 Tk	54	43.2
	15,001-20,000 Tk	22	17.6
	>20,000 TK	22	17.6
<b>Residential status</b>	Urban	57	45.6
	Rural	68	54.4
<b>Level of education</b>	Informal education	13	10.4
	Primary School	59	47.2
	Secondary School	37	29.6
	Above Secondary School	16	12.8
<b>Present Occupation</b>	Housewife	108	86.4
	Service holder	6	4.8
	Business	1	0.8
	Others	10	8.0
<b>Husband education</b>	Informal education	1	0.8
	Primary School	56	44.8
	Secondary School	44	35.2
	Above Secondary School	24	19.2
<b>Husband Occupation</b>	Service holder	24	19.2
	Farmer	14	11.2
	Business	34	27.2
	Others	53	42.4

**Figure 1** represents religion of the respondents where most of them 119(95.2%) were Muslims and only 6(5%) were Hindu.



**Figure 1: Distribution of respondents according to religion (n= 125)**

It can be shown from **figure 2** that 59.2% respondents were belongs to joint family and 40.8% were from nuclear family.



**Figure: 2 Distribution of respondents according to family type (n = 125)**

### Risk factors of eclampsia patients

**Table 2** shows the risk factors of eclampsia among the women. A total 125 respondents were interviewed, among them a large number of women were primipara 64(51.2%) and 61(48.8%) were multipara, gap between pregnancy was <1year among 7(5.6%) respondents, 1-2years among 32(25.6%) and >2years among 22(17.6%) women. A number of 100(80.0%) respondents had oedema and minority 25(20.0%) had no oedema. Majority of 82(65.6%) the respondents had no anemia and 43(34.4%) had anemia. Most of the respondents 119(95.2%) had no previous history of eclampsia and only 6(4.8%) had previous history of eclampsia. Among 125 respondent's majority of them 119(95.2%) had no diabetes, 113(90.4%) had no hypertension and only 1(0.8%) had renal disease.

**Table 2 Risk factors of eclampsia**

Risk factors of eclampsia		Frequency	% Distribution
<b>Parity</b>	Primi	64	51.2
	Multi	61	48.8
<b>Inter pregnancy space</b>	<1 year	7	5.6
	1-2 years	32	25.6
	>2 years	22	17.6
<b>History of multiple pregnancy</b>	Yes	4	3.2
	No	57	45.6
<b>Previous history of eclampsia</b>	Yes	6	4.8
	No	119	95.2
<b>Edema</b>	Yes	100	80.0
	No	25	20.0
<b>Anaemia</b>	Yes	43	34.4
	No	82	65.6
<b>Diabetes</b>	Yes	6	4.8
	No	119	95.2
<b>Hypertension</b>	Yes	12	9.6
	No	113	90.4
<b>Renal disease</b>	Yes	1	0.8
	No	124	99.2


**Maternal outcome of eclampsia patients**

**Table 3** showed maternal complications and outcome of eclampsia patients. Study findings revealed that (1.8%) eclampsia patients were developed pulmonary edema, 2.4% had HELP syndrome, 4.0% had obstetric shock and this study also found 0.8% mothers were died due to eclampsia.

**Table 3 Maternal complications of eclampsia patients (n=125)**

Complications	Frequency	Percent (%)
Disseminated Intravascular Coagulopathy (DIC)	0	0
Acute renal failure	0	0
Pulmonary edema	1	1.8
HELLP syndrome	3	2.4
Obstetric shock	5	4.0
Death	1	0.8
None	115	91.0
<b>Total</b>	<b>125</b>	<b>100</b>


Eclampsia patient’s occurrence of fits is shown in **figure 3**. Majority of the patients fits during their antepartum period (76.8%) [Figure-3].



**Figure 3: Distribution of the respondents according to occurrence of fits (n=125)**



**Figure 4** depicts mode of delivery among eclampsia patients. It can be seen that most of the (66%) respondents had caesarean section and 34% had normal vaginal delivery.



**Figure 4: Distribution of the respondents according to the mode of delivery**

### Neonatal outcomes of eclampsia patients

Neonatal outcomes of eclampsia patients are shown in **table 3**. 8.0% neonate were dead, 6.4% were stillborn and 12.8% were IUD. Birth asphyxia was present among 48.8% neonates, 66.4% neonates were low birth weight and 68.0% were preterm newborn. Among 125 mothers with eclampsia 28% neonates had Intra uterine growth retardation.

**Table 3 Neonatal outcomes of eclampsia patients**

Neonatal outcomes		Frequency	Percent (%)
<b>Perinatal outcome</b>	Alive	91	72.8
	Dead	10	8.0
	Still born	8	6.4
	IUD	16	12.8
<b>Birth Asphyxia (n=107)</b>	Present	61	48.8
	Absent	48	38.4
<b>Birth Weight</b>	Normal birth weight	42	33.6
	Low birth weight	83	66.4
<b>Maturity</b>	Preterm	85	68.0
	Full term	40	32.0
<b>IUGR (Intra uterine growth retardation)</b>	Yes	35	28.0
	No	90	72.0

### Discussion

After PPH, eclampsia is the second most frequent cause of maternal death. In Bangladesh, eclampsia is responsible for approximately 16% of maternal deaths. In the developing world, the mortality rate from hypertension diseases is substantially higher. Death in hypertensive disorders is associated with inadequate or absent prenatal treatment<sup>2,29</sup>. This study was designed to assess the maternal and fetal outcomes of eclampsia patients in Bangladesh.

In our study, most of the respondents (44.8%) were between age group 20-25 years. Many studies in Bangladesh found that eclampsia is more prevalent below 25 years of women<sup>24, 30</sup>. In our study, most of the respondents (44.8%) were between age group 20-25 years. Many studies in Bangladesh found that eclampsia is more prevalent below 25 years of women<sup>24, 30</sup>. This study showed most of the patients had poor economic condition (43.2%) that is consistent with another findings of the study that 52% of the

patients were belongs to poor and lower middle class<sup>[34, 16]</sup>, a great number of them had primary level of education (47.2%) which was also found similar with another study that 75% patients were either illiterate or got primary education only<sup>16, 33</sup>. This study also revealed most of the women were housewife (86.4%), most of them were primi gavida (51.2%) and majority of them comes from rural area (54.4%) which is in line with another study results<sup>30, 35, 24</sup>.

In our research 76.8% patients were fit during their antepartum period, Begum M R and Choudhary P also showed in their study that >75 % patients were in antepartum eclampsia group<sup>35, 24</sup>. In our study most of the (66%) respondents had caesarean section and 34% had normal vaginal delivery. Another study conducted in Bangladesh in 2014 found that among the antepartum eclamptic patients 90% had spontaneous vaginal delivery and only 10% had delivered by LUCS which indicates that nowadays caesarean delivery is increasing in Bangladesh<sup>32</sup>.

This study observed maternal complications due to eclampsia were pulmonary edema, HELLP syndrome and obstetric shock 0.8%, 2.4% and 4.0% respectively. Begum MR also found that pulmonary oedema and obstetric shock were 10.3 % and 7 % respectively<sup>24</sup>. Our study found only 0.8% deaths among eclampsia patients which is in contrast with another research findings that 8% mother were died due to eclampsia<sup>30</sup>. The difference of the findings may be due to early hospital arrival and availability of better treatment facilities.

Eclampsia not only kills mother but also kills fetus. It had been found from our research that 8.0% neonate were dead, 6.4% were stillborn and 12.8% were IUD. Similar with our findings in Shahabuddin's study they found that stillbirth was 24.50% and neonatal death was 8.40% among eclamptic patients<sup>15</sup>. Neonatal complications found in this research were birth asphyxia among 48.8% neonates, 66.4% neonates were low birth weight and 68.0% were preterm newborn. Among 125 mothers with eclampsia 28% neonates had Intra uterine growth retardation. Several studies showed perinatal deaths were higher in low-birth-weight babies<sup>14, 31</sup>. In shahabuddin's study, carried in Bangladesh, asphyxia was a major cause of neonatal loss with or without prematurity<sup>15</sup>. Irin's Study showed that the common causes of foetal deaths are birth asphyxia and prematurity<sup>31</sup>. Our study has significant limitations because we selected a smaller sample size due to time and financial constraints.

### **Conclusion and recommendations**

This study found that pulmonary edema, HELLP syndrome, obstetric shock were most common complications of eclampsia and majority of the patients fits during their antepartum period. Perinatal death rate remains high in eclamptic women. Still birth, prematurity, IUGR and birth asphyxia were found as important causes of perinatal loss among the eclampsia patients. Early diagnosis and management of pre-eclampsia and hypertensive diseases are important to prevent eclampsia. Early referral of eclampsia patients, better obstetric management, early resuscitative measures and good neonatal care facilities can improve the perinatal outcome.

## References

1. Damien, S, Patgric, G, Francies, P, Pierre, L, Serge, B, Gerad, B, et al. Aspirin (100mg) used for prevention of preeclampsia in nulliparous women, the Eassai Regional Aspirin Mere – Enfant Study (Part-1). *Br J Obstet Gynaecol* 2003; 110: 475-84.
2. Philip JS, Mark PL, Tina KJ, Chapple. Maternal blood pressure in pregnancy, birth weight and perinatal mortality is first birth. *BMJ* 2004; 329: 1312-4.
3. Amaral LM, Wallace K, Owens M, La Marca B. Pathophysiology and Current Clinical Management of Preeclampsia. *Curr Hypertens Rep.* 2017 Aug; 19(8):61. doi: 10.1007/s11906-017-0757-7. PMID: 28689331; PMCID: PMC5916784.
4. Ross MG, Meyer BA, Telavera F, Ramus RM. Eclampsia Overview. *Medscape* 253960. 2011:1–13.
5. Ade-Ojo IP, Loto OM. Outcome of eclampsia in OAUTH Ile-Ife. *Nig J Clin Pract.* 2008; 11(3):279–284.
6. Douglas KA, Redman CW. Eclampsia in the united kingdom *BMJ* 1994; 309:1395- 400. DOI: 10.1136/bmj.309.6966.1395.
7. Crowther CA. Eclampsia at Harare Maternity Hospital. An epidemiological study. *S Afr Med J.* 2009; 68(13):927-929.
8. Bergström S, Povey G, Songane F, Ching C. Seasonal incidence of eclampsia and its relationship to meteorological data in Mozambique. *J Perinat Med.* 2012; 20(2):153-158. DOI: 10.1515/jpme. 2012.20.2.153.
9. Mahler H. The safe motherhood initiative: A call to action. *Lancet.* 2011; 1(8534):668- 670. DOI: 10.1016/s0140-6736(87)90423-5.
10. Rosenfield A, Maine D. Maternal mortality a neglected tragedy. Where is the M in MCH? *Lancet* 2005; 2(8446):83-5. Available:[https://doi.org/10.1016/S0140-6736\(85\)90188-6](https://doi.org/10.1016/S0140-6736(85)90188-6).
11. Duley L. Maternal mortality associated with hypertensive disorders of pregnancy in Africa, Asia, Latin America and the Caribbean. *Br J Obstet Gynecol* 2012; 99(11):937.
12. Fauveau V, KNij KA, Chakrabarty J and Chowdhury AL. Causes of maternal mortality in rural Bangladesh. *Bull WHO.* 2008; 66:643-651.
13. Niport, Mitra and associate, ORC macro, Bangladesh republic and health survey, 2004; 123.
14. Shamsuddin L, Rouf S, Khatoun H. Perinatal outcome in Eclampsia. *Bangladesh J Obstet Gynaecol* 2005; 10:65-72. DOI: 10.36347/sjams.2020.v08i09.024.
15. Shahabuddin AKM, Hasnat M, Hamid T, Rahman AKMF. Perinatal outcome in Eclampsia. *Bangladesh J child Health.* 2016; 20; 8-14.
16. Shabnam Rahman, Kamrun Nessa, Khaled Kaiser, and Runa Laila, Fetomaternal Outcome of Eclampsia in Shaheed Suhrawardy Medical College and Hospital, Dhaka, Bangladesh; *Asian Research Journal of Gynaecology and Obstetrics*5(1): 49-57, 2022;Article no.ARJGO.85786.
17. Nahar S. Utility of misoprostol for labour induction in severe preeclampsia and eclampsia .*The Journal of Obstetrics & Gynecology Reserch.* 2004; 30:349-352.
18. Bangladesh Maternal Mortality and Health Care Survey 2016.
19. Datta DC. *Textbook of obstetric including perinatilology and contraception.*4th ed. Calcutta.1998; 51, 51, 236-9, 648-9.
20. Guerrier et al, Factors associated with severe preeclampsia and eclampsia in Jahun, Nigeria *International Journal of Women's Health* 2013;5 509–513.
21. Ndaboine et al *Maternal and Perinatal Outcomes among Eclamptic Patients Admitted to Bugando Medical Centre, Mwanza, Tanzania African Journal of Reproductive Health* March 2012; 16(1): 36.
22. Rayamajhi A K\*, Uprety D\*, Agrawal A\*, Pokhrel H\* *FETOMATERNAL OUTCOME IN ECLAMPSIA Journal of Nepal Medical Association* 2003; 42: 341-345.
23. Sultana, Kanij and Amy Dempsey. 2017. "Knowledge of pre-eclampsia and eclampsia in Bangladesh," *Ending Eclampsia Brief.* Washington, DC: Population Council.
24. Tahmina HZ, Shahid AR, Hosna AU, Alam A "Study on Outcome of Eclampsia Patients in District Hospital in Bangladesh" *J Dhaka Med Coll.* 2014; 23(2) : 223-226.
25. Villar MA, Sibai BM. Eclampsia. *Obset Gynaecol North Am.* 1988; 15: 355-377.13.
26. Sibai BM, Villar MA, Mabie BC. Acute renal failure in hypertensive disorders of pregnancy. Pregnancy outcome and remote prognosis in thirty-one consecutive cases. *Am J Obstet Gynecol.* 1990; 162(3):777-783.
27. Connell H, Dalgleish JG, Downing JW. General anaesthesia in mothers with severe pre-eclampsia/eclampsia. *Br J Anaesth.* 1987; 59(11):1375-1380.
28. Onuh SO, Aisien AO. Maternal and fetal outcome in eclamptic patients in Benin City, Nigeria. *J Obstet Gynaecol.* 2004; 24(7):765-768.
29. Reynold C, Mabie WC, Sibai BM, Decherny AH, Nathan, L. *Hypertensive status of pregnancy.* 9th ed. New York: McGraw-Hill; 2003: p.338-53.
30. Akhtar, R., Ferdous, A., & Bhuiyan, S. N. (2013). Maternal and Fetal Outcome of Eclamptic Patients in a Tertiary Hospital. *Bangladesh Journal of Obstetrics & Gynaecology*, 26(2), 77–80. <https://doi.org/10.3329/bjog.v26i2.13784>.

31. Irin Parveen Alam, Sayeba Akhter. *Perinatal outcome of eclampsia in Dhaka Medical College Hospital. Bangladesh J Obstet Gynecol.* 2008; 23(1):20-24.
32. Tahmina, H. Z., Shahid, A. R., Hosna, A. U., & Alam, A. (2014). *Study on Outcome of Eclampsia Patients in District Hospital in Bangladesh. Journal of Dhaka Medical College,* 23(2), 223–226. <https://doi.org/10.3329/jdmc.v23i2.25395>.
33. Bugalho A, Bacci A, Bergstrom S. *Risk factors in Mozambican women with Eclampsia: Afr J Reprod Health.* 2001 Aug; 5(2)30-5. DOI: 10.2307/3583428 20.
34. Dare FO, Eniola OA, Bariweni AC. *Eclampsia revisited. Nig J Med.* 1998; 7: 168-171.
35. Choudhary P. *Eclampsia: A hospital based retrospective study. Kathmandu University Medical Journal .*2003; 1: 237-241.
36. Mou, A.D., Barman, Z., Hasan, M. et al. *Prevalence of preeclampsia and the associated risk factors among pregnant women in Bangladesh. Sci Rep* 11, 21339 (2021). <https://doi.org/10.1038/s41598-021-00839>.